Table S3. Association of NF90 expression with patients' clinicopathological features in ovarian carcinomas

	NF90 Expression			
Feature	All cases	Low	High	P value ^a
Age [□] (years)				0.079
≤ 50.8	83	43 (51.8%)	40 (48.2%)	
> 50.8	86	56 (65.1%)	30 (34.9%)	
Histological type				0.26
serous	113	70 (61.9%)	43 (38.1%)	
mucinous	21	9 (42.9%)	12 (57.1%)	
Others ^c	35	20 (57.1%)	15 (42.9%)	
Histological grade ^d				0.015
G1	29	10 (34.5%)	19 (65.5%)	
G2	100	63 (63.0%)	37 (37.0%)	
G3	40	26 (65.0%)	14 (35.0%)	
pT status				0.155
pT1	47	22 (46.8%)	25 (53.2%)	
pT2	32	20 (62.5%)	12 (37.5%)	
pT3	90	57 (63.3%)	33 (36.7%)	
pN status				0.017
pN0	83	41 (49.4%)	42 (50.6%)	
pN1	86	58 (67.4%)	28 (32.6%)	
pM status				0.001
рМХ	146	78 (53.4%)	68 (46.6%)	
pM1	23	21 (91.3%)	2 (8.7%)	
FIGO ^e stage				<0.001
I	30	11 (36.7%)	19 (63.3%)	
II	20	9 (45.0%)	11 (55.0%)	
III	96	58 (60.4%)	38 (39.6%)	
IV	23	21 (91.3%)	2 (8.7%)	
DICER expression				<0.001
Low expression	69	59 (85.5%)	10 (14.5%)	
High expression	100	40 (40.0%)	60 (60.0%)	

^aChi-square test; ^bmean age at surgery; ^cincludes endometrioid, clear cell and undifferentiated carcinoma; ^dSilveberg classification; ^eInternational Federation of Gynecology and Obstetrics